



VARIANT OF LAX-MILGRAM LEMMA FOR BANACH SPACES

Gheorghe Doboș

To Professor Silviu Sburlan, at his 60's anniversary

Abstract

In the paper, it is shown that, by taking a condition on the difference $|a(u, v) - a(v, u)|$ in the Lax-Milgram Lemma, we may apply it to the same problem in a Banach space.

We recall the **Lax-Milgram Lemma**:

Let X be a real Hilbert space, let $a(\cdot, \cdot)$ be a continuous, coercitive, bilinear form defined on X , and let f be a continuous linear form on X . Then there exists one and only one element $u \in X$ which satisfies

$$(\forall)v \in X, a(u, v) = f(v).$$

We establish the conditions of our work.

Let X be a Banach space, $a : X \times X \rightarrow \mathbb{R}$ be a bilinear form and $f : X \rightarrow \mathbb{R}$ be a linear continuous function.

Assume that:

- 1) $\exists M > 0$ a.i. $|a(u, v)| \leq M \|u\| \cdot \|v\|$ (boundness);
- 2) $\exists K^2 > 0$ a.i. $a(u, u) \geq K^2 \|u\|^2$ (coercitivity).

Let

$$\alpha = \sup_{u, v \neq 0} \frac{|a(u, v) - a(v, u)|}{\|u\| \|v\|}$$

From $|a(u, v)| \leq M \|u\| \cdot \|v\|$ we get $|a(u, v) - a(v, u)| \leq 2M \|u\| \|v\|$ and therefore there exists α and $\alpha \leq 2M$.

We denote $b(u, v) = a(u, v) + a(v, u)$ and for all $x \in X$ we denote $g_x(v) = f(v) + a(v, x)$

Obviously b becomes a bilinear form which is symmetric, continuous, and coercitive on X and g_x is linear and continuous.

Denoting

$$J_x(u) = \frac{1}{2}b(u, u) - g_x(u) = a(u, u) - g_x(u),$$

and using Ritz method, there exists a unique $u_x \in X$ which satisfies $J_x(u_x) = \inf_{u \in X} J_x(u)$ and in addition satisfies a variational equation

$$b(u_x v) = g_x(v), \quad (\forall) v \in X \text{ i.e.}$$

$$a(u_x, v) + a(v, u_x) = f(v) + a(v, x), \quad (\forall) v \in X$$

By starting with $u_0 \in X$, we define the sequence $(u_n)_{n \in \mathbb{N}}$ given by

$$a(u_n, v) + a(v, u_n) = f(v) + a(v, u_{n-1}).$$

We prove that:

$$\|u_n - u_{n-1}\| \leq \left(\frac{\alpha}{4K^2} + \frac{1}{2}\right) \|u_{n-1} - u_{n-2}\|, \quad n \geq 2,$$

where α and K are the nonsymmetry constant and, respectively the coercitivity constant for a .

Remark 1. The following inequality holds:

$$\begin{aligned} a\left(u_n - u_{n-1} - \frac{1}{2}(u_{n-1} - u_{n-2}), u_n - u_{n-1} - \frac{1}{2}(u_{n-1} - u_{n-2})\right) &\geq \\ &\geq K^2 \left\|u_n - u_{n-1} - \frac{1}{2}(u_{n-1} - u_{n-2})\right\|^2. \end{aligned}$$

From $a(u_n, v) + a(v, u_n) - f(v) - a(v, u_{n-1}) = 0$, and $a(u_{n-1}, v) + a(v, u_{n-1}) - f(v) - a(v, u_{n-2}) = 0$,

we get

$$a(u_n - u_{n-1}, v) + a(v, u_n - u_{n-1}) = a(v, u_{n-1} - u_{n-2}).$$

Denoting $v_n = u_n - u_{n-1}$, $n \geq 1$, we have

$$\begin{aligned} a(v_n, v) + a(v, v_n) &= a(v, v_{n-1}) = \frac{1}{2}(a(v, v_{n-1}) + a(v_{n-1}, v)) + \\ &\quad + \frac{1}{2}(a(v, v_{n-1}) - a(v_{n-1}, v)). \end{aligned}$$

We deduce immediately:

$$\begin{aligned} a\left(v_n - \frac{1}{2}v_{n-1}, v\right) + a\left(v, v_n - \frac{1}{2}v_{n-1}\right) &= \frac{1}{2}[a(v, v_{n-1}) - a(v_{n-1}, v)], \\ a\left(v_n - \frac{1}{2}v_{n-1}, v_n - \frac{1}{2}v_{n-1}\right) &= \\ = \frac{1}{4}\left[a\left(v_n - \frac{1}{2}v_{n-1}, v_{n-1}\right) - a\left(v_{n-1}, v_n - \frac{1}{2}v_{n-1}\right)\right] &\leq \\ \leq \frac{\alpha}{4}\|v_{n-1}\| \cdot \left\|v_n - \frac{1}{2}v_{n-1}\right\| \end{aligned}$$

and, from Remark 1, we get the inequalities:

$$\begin{aligned} K^2 \left\|v_n - \frac{1}{2}v_{n-1}\right\| &\leq \frac{\alpha}{4}\|v_{n-1}\| \cdot \left\|v_n - \frac{1}{2}v_{n-1}\right\|, \\ \left\|v_n - \frac{1}{2}v_{n-1}\right\| &\leq \frac{\alpha}{4K^2}\|v_{n-1}\|, \\ \|v_n\| &\leq \left\|v_n - \frac{1}{2}v_{n-1}\right\| + \frac{1}{2}\|v_{n-1}\| \leq \left(\frac{\alpha}{4K^2} + \frac{1}{2}\right)\|v_{n-1}\|. \end{aligned}$$

Proposition 2. *If $\Lambda = \frac{\alpha}{4K^2} + \frac{1}{2} < 1$, for $\alpha < 2K^2$, the sequence $(X_n)_n$ is convergent.*

Proof. We give the calculation for showing that $(u_n)_{n \in \mathbb{N}}$ is a fundamental sequence in the Banach space. First, we have:

$$\|u_n - u_{n-1}\| \leq \Lambda \|u_{n-1} - u_{n-2}\| \leq \dots \leq \Lambda^{n-1} \|u_1 - u_0\|;$$

therefore

$$\begin{aligned} \|u_{n+p} - u_n\| &\leq \|u_{n+p} - u_{n+p-1}\| + \dots + \|u_{n+1} - u_n\| \leq \\ &\leq (\Lambda^{n+p-1} + \dots + \Lambda^n) \|u_1 - u_0\| \leq \frac{\Lambda^n}{1 - \Lambda} \|u_1 - u_0\|, \quad \Lambda \in (0, 1) \end{aligned}$$

and $(u_n)_n$ is fundamental

As X is complete, it follows that $(u_n)_n$ is convergent.

Let $u^* = \lim u_n$. As $a(u_n, v) + a(v, u_n) = f(v) + a(v, u_{n-1})$ we have $a(u^*, v) = f(v)$, $(\forall) v \in V$, therefore, by weakening the nonsymmetry of the bilinear form, we may extend the Lax-Milgram Lemma to a Banach space. In addition, the solution is the limit of a recurrence sequence obtained by minimizing the energy functional J .

References

- [1] Ciarlet Ph., *Numerical analysis of the finite element method*, Les Presses de L'Université de Montreal, 1976.
- [2] Dincă Gh., *Metode variaționale și aplicații*, Ed. Tehnică, 1980.

"Ovidius" University of Constantza,
Faculty of Mathematics and Informatics,
Mamaia Bd., 124,
8700 Constantza,
Romania